

The Relationship of *In Vitro* Anti-Inflammatory and Antioxidant Activity with Total Phenolic and Flavonoids Content of *Tithonia diversifolia* Leaves Ethanolic Extract and Fractions

Endah Puspitasari*, Nuri, Siti Muslichah, Bawon Triatmoko, Dewi Dianasari

Bioactive Natural Product Development Research Group, Faculty of Pharmacy, University of Jember, Jember, Indonesia

*Corresponding author: e.puspitasari@unej.ac.id

Received: January 9, 2024; **Accepted:** November 4, 2024; **Published:** November 26, 2024

Abstract

A previous study showed that *Tithonia diversifolia* ethanolic extract has antiproliferative properties against HeLa cervical cancer cells. It has also shown anti-inflammatory properties *in vivo*, one of the approaches to cancer chemoprevention. This research aimed to determine the relation of *in vitro* anti-inflammatory and antioxidant activity of *T. diversifolia* leaves ethanolic extract and fractions with total phenolic and flavonoid content. The anti-inflammatory was evaluated *in vitro* using red blood cell (RBC) membrane stabilization and inhibition of protein denaturation method, while antioxidant activity was done using the DPPH method. Phytochemical screening was done using the TLC method. Total phenolic and flavonoid content were also counted using the colorimetry method. ANOVA followed by LSD post hoc, multiple regression analysis, and Pearson correlation were used to analyze the data. Compared to its fractions, the highest anti-inflammatory and antioxidant activity was obtained from *T. diversifolia* leaves ethanolic extract. The *T. diversifolia* leaves ethanolic extract contained terpenoids, alkaloids, polyphenols, and flavonoids. Overall, the antioxidant property contributed more to RBC membrane stabilization than protein denaturation inhibition. The phenolic and flavonoid content contributed most to antioxidant and red blood cell membrane stabilization rather than protein denaturation inhibition.

Keywords: Anti-inflammatory; Antioxidant; Phytochemical screening; *Tithonia diversifolia* leaves extract and fractions; Total phenolic and flavonoids content

1. INTRODUCTION

Inflammation is a defense mechanism to local injury that protects from tissue damage caused by physical, chemical, or microbiological trauma (Tu et al., 2022). Inflammation is the cause of many diseases, including cancer (Greten & Grivennikov, 2019). The incidence of cancer in the world is increasing. According to estimates (Siegel et al., 2023), there were 609,820 cancer deaths and 1,958,310 new cancer cases in the US alone. Cancer is the cause of one death out of every six (World Health Organization, 2022). An approach to cancer chemoprevention is anti-inflammatory drugs (Greten & Grivennikov, 2019). First-line anti-inflammatory drugs can cause serious side effects, e.g., impaired renal function, hypertension, edema, gastrointestinal bleeding, and even increase the risk of heart failure, stroke, myocardial infarction, and death (Bindu et al., 2020). To avoid these side effects caused by anti-

inflammatory drugs, researchers are exploring new anti-inflammatory candidates, including those from natural products, e.g., plants (Nunes et al., 2020). One candidate that could be explored as an anti-inflammatory agent is *Tithonia diversifolia* (Hemsl) A. Gray.

A previous study showed that ethanolic extract of *T. diversifolia* leaves exhibited antiproliferative activity on HeLa cervical cancer cells (Puspitasari et al., 2022). *T. diversifolia* is used as an anti-inflammatory preparation by society (Dalimartha, 1999). Ethanolic extract of *T. diversifolia* leaves exhibited anti-inflammatory activity *in vivo* by reducing the exudate volume of carrageenan-induced mice paw edema models (Lubis et al., 2020). The polar extract of *T. diversifolia* leaves contains chlorogenic acid and sesquiterpene lactone, which show better anti-inflammatory effects than indomethacin (Lopes et al., 2021). *T. diversifolia* leaves exhibit antioxidant activity (Tamfu et al., 2022). Compounds with antioxidant activity usually have anti-inflammatory properties (Bourais et al., 2023). This research was conducted to reveal a detailed mechanism of action of the anti-inflammatory properties of *T. diversifolia* leaves ethanolic extract and their fractions, i.e., n-hexane, dichloromethane, and methanol-water fraction, their antioxidant properties, as well as to determine the relationship of anti-inflammatory and antioxidant activity with total phenolic and flavonoids content. Fractionation of an extract is a further step for natural products' isolation (Xin et al., 2021). Thus, this research was aimed at preliminary bioactivity-guided isolation for the anti-inflammatory and antioxidant activity of *T. diversifolia* leaves ethanolic extract.

2. MATERIAL AND METHODS

2.1. Extraction and fractionation of *T. diversifolia* leaves

T. diversifolia leaves were collected from Jember. They were hand-picked based on healthy and mature leaves and taken from a mature plant starting from the 5th strand from the tip of the stem to the base. The plant identity was confirmed using plant determination at the Botany Laboratory, Department of Biology, Faculty of Mathematics and Natural Sciences University of Jember. The leaves were air-dried, pulverized, and sifted prior to extraction. The extraction used maceration on 100-gram leaves with continuous stirring with 96% ethanol as solvent (1:7 ratio). The maceration was repeated three times. The extract was evaporated using a rotary evaporator (Heidolph Laborota 4000) until a thick extract was obtained (Puspitasari et al., 2022). A 20-gram thick ethanolic extract was then diluted with methanol-water (1:1), followed by fractionation using the liquid-liquid partition method with n-hexane, dichloromethane, and methanol-water using a 1:1 ratio. The fractions were evaporated using a rotary evaporator until thick fractions were obtained.

2.2. Anti-inflammatory assay using red blood cells membrane stabilization method

Ethical clearance was obtained from KEPK FKG Jember University with number 1423/UN25.8/KEPK/DL/2021. The red blood cell (RBC) membrane stabilization method was done based on (Yesmin et al., 2020) with slight modification. RBCs were obtained from female rabbits, placed in EDTA tubes, and centrifuged at 3,000 rpm for 15 min at room temperature.

The residue was washed with isosaline until the supernatant was clear. The RBC obtained was suspended 10% in isosaline.

All of the experiments were carried out in three replicates. A test solution of 4.5 mL was prepared by adding 1 mL of 0.15 M PBS pH 7.4 with 0.5 mL of RBC suspension (2% v/v), 1 mL of sample solution with the series concentration of 500, 250, 125, 62.5, and 31.3 µg/mL, and 2 mL of hyposaline solution (0.25% b/v). Sodium diclofenac (Phapros, series concentration of 500, 250, 125, 62.5, and 31.3 µg/mL) was used as a positive control. The mixture was incubated at 56 °C for 30 min in a water bath, followed by centrifugation at 5,000 rpm for 10 min at room temperature, and read at 560 nm absorbance using an ELISA reader (Dialab EL X800). The stability of red blood cells was then calculated based on Formula 1.

$$\text{Red blood cell membrane stability (\%)} = \frac{(100 - (\text{sample absorbance} - \text{control absorbance}))}{(\text{control absorbance})} \times 100\% \quad [1]$$

Formula 1. The calculation of the percentage stability of red blood cell membrane.

The plot of concentration vs. RBC membrane stability was made as the basis of the IC₅₀ calculation.

2.3. Anti-inflammatory assay using protein denaturation inhibition method

The protocol used for this method was adapted from Saleem et al. (2020). The reaction mixture consisted of 0.2 mL of 1% bovine serum albumin (BSA, BioPLUS) and 2.8 mL of PBS (pH adjusted to 6.4 with 1N HCl) to which normal saline (negative control), test extract/fractions (500, 250, 125, 62.5, and 31.3 µg/mL), or sodium diclofenac (500, 250, 125, 62.5, and 31.3 µg/mL) were added as positive controls. The mixture was incubated at 37 °C for 20 minutes and heated to 51 °C for 20 minutes. After cooling, the absorbance was measured at 660 nm. The experiment was carried out in three replicates. Inhibition of protein denaturation was calculated based on Formula 2.

$$\text{Protein denaturation inhibition (\%)} = \frac{(\text{control absorbance} - \text{sample absorbance})}{(\text{control absorbance})} \times 100\% \quad [2]$$

Formula 2. The calculation of the percentage protein denaturation.

The plot of concentration vs protein denaturation inhibition was made as the basis of the IC₅₀ calculation.

2.4. Antioxidant assay using DPPH method

The antioxidant assay was done using the DPPH microplate assay. Briefly, the extract and fractions were diluted using methanol (Merck), and then 150 µl of the sample was placed in a 96-well plate with 100 µl of 0.2 mM DPPH solution (Sigma Aldrich). After incubation for 30 minutes at room temperature in a dark room, the absorbance was measured at 515 nm using an ELISA reader (Dialab EL X800). Vitamin C (Emsure, concentration of 21.2, 10.6, 5.3, 2.65, 1.325, 0.663, 0.331, and 0.116 µg/mL) was used as a positive control. The experiment was carried out in three replicates. A series of concentrations used ranging from 100.6-0.686 µg/mL for the ethanolic extract, 500-3.906 µg/mL for n-hexane fraction, 106-0.828 µg/mL for

dichloromethane fraction, and 104-0.813 $\mu\text{g}/\text{mL}$ for methanol-water fraction, to make sure that the IC_{50} calculated by interpolation of each sample. The antioxidant activity was calculated based on Formula 3.

$$\text{DPPH scavenging activity (\%)} = \frac{(\text{control absorbance} - \text{sample absorbance})}{(\text{control absorbance})} \times 100\% \quad [3]$$

Formula 3. The calculation of percentage antioxidant activity by DPPH scavenging activity.

The plot of concentration vs DPPH scavenging activity was made as the basis of the IC_{50} calculation (Nasution et al., 2019).

2.5. Phytochemical screening

The phytochemical screening of the extract and fractions of *T. diversifolia* leaves was done based on the TLC method with the respective chemical spray to detect terpenoids (anisaldehyde sulfuric acid), alkaloids (Dragendorff's reagent), polyphenols (FeCl_3), and flavonoids (boric citrate) (Moulishankar, 2021). The TLC was done using silica gel 60 F₂₅₄ as the stationary phase. In contrast, the mobile phase was the optimum composition, resulting in the best separation and resolution for the spots found in each sample. The mobile phase used was and n-hexane (Merck): ethyl acetate (Merck) (3:2) for ethanol extract, toluene (Merck): ethyl acetate (Merck) (7:3) for n-hexane fraction, n-hexane (Merck): ethyl acetate (Merck) (2:1) for dichloromethane fraction, and n-hexane (Merck): ethyl acetate (Merck): methanol (Merck) (1:1:2) for methanol-water fraction.

2.6 Total phenolic content

The experiment was carried out in three replicates. A 100 μl sample (extract or fractions at the concentration of 1,060 $\mu\text{g}/\text{mL}$ (ethanolic extract); 1,010 $\mu\text{g}/\text{mL}$ (n-hexane fraction); 1,040 $\mu\text{g}/\text{mL}$ (dichloromethane fraction); and 1,000 $\mu\text{g}/\text{mL}$ (methanol-water fraction)) solution in methanol (Merck) was reacted with 100 μl Folin Ciocalteu reagent (Merck) in 96 healthy plate. A 50 μl of Na_2CO_3 (Merck) was added. After 30 min of incubation at room temperature in a dark place, the absorbance was measured at 735 nm using an ELISA reader (Dialab EL X800). The phenol content was calculated by plotting the sample absorbance into the standard curve of gallic acid (Sigma Aldrich, concentration of 50, 40, 30, 20, and 10 $\mu\text{g}/\text{mL}$) (Loucif et al., 2020).

2.7. Total flavonoid content

The experiment was carried out in three replicates. Sample (extract or fractions at the concentration of 1,060 $\mu\text{g}/\text{mL}$ (ethanolic extract); 1,010 $\mu\text{g}/\text{mL}$ (n-hexane fraction); 1,040 $\mu\text{g}/\text{mL}$ (dichloromethane fraction); and 1,000 $\mu\text{g}/\text{mL}$ (methanol-water fraction)) was dissolved in methanol and pipetted 100 μl in a 96 well plate, followed by the addition of 100 μl AlCl_3 50 μl CH_3COOK 10%, and 50 μl methanol (Merck). The incubation was done for 30 min in a dark room at room temperature. Then, the absorbance was measured at 425 nm using an ELISA reader (Dialab EL X800). The flavonoid content was calculated as quercetin equivalent based

on a quercetin (Sigma Aldrich) standard curve plot with concentrations of 40, 30, 25, 20, and 10 $\mu\text{g/mL}$ (Loucif et al., 2020).

2.8. Data analysis

The IC_{50} obtained from the anti-inflammatory and antioxidant assay and the total phenolic and flavonoid content were evaluated for their differences using ANOVA, followed by LSD using a 99% confidence level. The multiple regression analysis was done to analyze whether total phenolic and flavonoid content contributed to the anti-inflammatory and antioxidant activity, while the Pearson correlation analysis was done to determine the correlation between parameters in this research.

3. RESULTS AND DISCUSSION

Plant determination has established that the plant's identity was *T. diversifolia*. The extract and fraction yields obtained are shown in Table 1. The yield difference was suggested due to the different solubility of the compounds fractionated by different solvent polarities. The highest yield was obtained from the methanol-water fraction, suggesting that the ethanolic extract of *T. diversifolia* leaves contained more polar compounds than non-polar and semi-polar compounds.

The anti-inflammatory activity of each sample is shown in Table 2. Even though the anti-inflammatory activity of the extract and fractions was incomparable to sodium diclofenac as the positive control for these assays, the IC_{50} for the extract and fractions was considered promising. All the samples exhibited significantly different anti-inflammatory in both methods. The ethanolic extract of *T. diversifolia* leaves showed the highest anti-inflammatory activity on both methods.

Table 1. Extract and fractions yield obtained from *T. diversifolia* leaves

Sample	Yield (%)
Ethanolic extract of <i>T. Diversifolia</i> leaves	23.425
N-hexane fraction of <i>T. Diversifolia</i> leaves	10.217
Dichloromethane fraction of <i>T. Diversifolia</i> leaves	22.751
Methanol-water fraction of <i>T. Diversifolia</i> leaves	49.215

The antioxidant activity of *T. diversifolia* leaves ethanolic extract and its fractions are shown in Table 3. The vitamin C as a positive control showed that the method had been carried out correctly. The IC_{50} values of all samples were significantly different. Both ethanolic extract and methanol-water fraction of *T. diversifolia* leaves showed strong antioxidant activity, but the other two (n-hexane and dichloromethane fraction of *T. diversifolia* leaves) exhibited intermediate antioxidant activity. Again, the ethanolic extract of *T. diversifolia* leaves showed the highest antioxidant activity.

The phytochemical screening showed that the ethanolic extract of *T. diversifolia* leaves contains more complex compounds compared to its fractions (Figure 1 and Table 4). The

fractionation did separate the compounds based on their polarity. Alkaloids did not appear in any fraction, suggesting that this compound might be too low to be detected by the TLC spray reagent to be observed with the eyes. Since alkaloids were found in the extract, they must have been found in any fractions. The sensitivity is the limitation of TLC spray reagents (Zhang et al., 2021). It might be found in the fractions if we used the colorimetric method using a spectrophotometer since instrumental analysis would be more precise. The instrumental analysis will provide more sensitivity, resulting in more reliable data (Zhang et al., 2021). Alkaloids in *T. diversifolia* leaves seem to be the lowest compounds found compared to others, e.g., terpenoids, polyphenols, and flavonoids. Studies showed that only two alkaloids are found in *T. diversifolia* leaves (Zhao et al., 2012).

Table 2. Anti-inflammatory activity of *T. diversifolia* leaves ethanolic extract and its fractions. *Description:* *IC₅₀ data was shown as mean \pm SD (n=3); different letter notations in the same column indicate significant differences (LSD, p<0.01) in red Blood Cells (RBC).

Sample	IC ₅₀ (μ g/mL)*	
	RBC membrane stabilization	Protein denaturation inhibition
Sodium diclofenac	60.315 \pm 0.729 ^a	2.171 \pm 0.025 ^f
Ethanolic extract of <i>T. diversifolia</i> leaves	78.427 \pm 0.594 ^b	2.529 \pm 0.044 ^g
n-hexane fraction of <i>T. diversifolia</i> leaves	94.500 \pm 0.206 ^c	3.745 \pm 0.085 ^h
Dichloromethane fraction of <i>T. diversifolia</i> leaves	89.268 \pm 0.224 ^d	2.937 \pm 0.036 ⁱ
Methanol-water fraction of <i>T. diversifolia</i> leaves	92.236 \pm 0.570 ^e	3.502 \pm 0.021 ^j

The ethanolic extract of *T. diversifolia* leaves and its fractions were then evaluated for their phenolic and flavonoid content. The total phenolic and flavonoid content is shown in Table 5. All the phenolic and flavonoid content were significantly different among the samples. The ethanolic extract of *T. diversifolia* leaves contained the highest phenolic compound. The ethanolic extract is more complex than the fractions, suggesting that the compounds in the ethanolic extract were being divided further into fractions (López-Cabeza et al., 2024). Thus, the ethanolic extract contained the highest phenolic compounds compared to its fractions.

Table 3. Antioxidant properties of *T. diversifolia* leaves ethanolic extract and its fractions. *Description:* *IC₅₀ data was shown as mean \pm SD (n=3), Different letter notations indicate significant differences (LSD, p<0.01), **DPPH antioxidant activity classification based on Phongpaichit et al. (2007).

Sample	IC ₅₀ ((μ g/mL)*	Antioxidant category**
Vitamin C	2.534 \pm 0.118	Very strongly active
Ethanolic extract of <i>T. diversifolia</i> leaves	22.592 \pm 0.468 ^a	Strongly active
n-hexane fraction of <i>T. diversifolia</i> leaves	67.755 \pm 0.674 ^b	Moderately active
Dichloromethane fraction of <i>T. diversifolia</i> leaves	51.105 \pm 0.805 ^c	Moderately active
Methanol-water fraction of <i>T. diversifolia</i> leaves	27.218 \pm 0.848 ^d	Strongly active

On the other hand, the highest flavonoid content was measured in the n-hexane fraction of *T. diversifolia* leaves. Some flavonoid compounds are non-polar polymethoxy aglycones or

isoflavones, in which sugar groups or glycoside forms are released (Satria et al., 2022). However, there is a possibility that the total flavonoid content obtained does not show the actual results. It is because extracts and fractions already have a color before being reacted; for example, the n-hexane fraction is green because it contains a lot of chlorophyll. Chlorophyll will be detected at a wavelength of 320-430 nm (Christiana et al., 2008), while the wavelength used for flavonoid detection with AlCl_3 is at a wavelength of 425 nm. At the test wavelength, chlorophyll is absorbed, so the calculated flavonoid content is above the actual value.




Figure 1. Chromatogram of phytochemical screening on ethanolic extract (A), n-hexane fraction (B), dichloromethane fraction (C), and methanol-water fraction of *T. diversifolia* leaves. The TLC was done using silica gel 60 F₂₅₄ as the stationary phase. The mobile phase used was n-hexane (Merck): ethyl acetate (Merck) (3:2) for ethanol extract (A); toluene (Merck): ethyl acetate (Merck) (7:3) for n-hexane fraction (B); n-hexane (Merck): ethyl acetate (Merck) (2:1) for dichloromethane fraction (C); and n-hexane (Merck): ethyl acetate (Merck): methanol (Merck) (1:1:2) for methanol-water fraction (D). The chromatogram sprayed using respective spray reagents for specific compounds, i.e., anisaldehyde sulfuric acid for terpenoids, characterized by purplish spots (a); Dragendorff's reagent for alkaloids, characterized by orangish spots (b); FeCl_3 for polyphenols, characterized by blackish spots (c); and boric citrate for flavonoids, characterized by yellowish spots which disappeared quickly (d). Smear color will lead to undetectable compounds visually (Zhang et al., 2021).

Some fractions, i.e., n-hexane and dichloromethane fractions of *T. diversifolia* leaves, resulted in higher flavonoid content than phenolic content. Theoretically, flavonoids are part of phenolic compounds. It is unlikely to have a higher flavonoid quantification than phenolic compounds. However, sometimes, the researchers found the same anomaly, the flavonoid content they had was higher than the phenolic compound (Yu et al., 2021; Zulkifli et al., 2020). Nevertheless, there needs to be a brief explanation of this phenomenon. Not all phenolic compounds could be determined by the Folin Ciocalteu reagent or simply because the reference substances used as standard are different.

Table 4. Phytochemical screening of *T. diversifolia* leaves ethanolic extract and its fractions.

Sample	Terpenoids	Alkaloids	Polyphenols	Flavonoids
Ethanolic extract of <i>T. diversifolia</i> leaves	+	+	+	+
N-hexane fraction of <i>T. diversifolia</i> leaves.	+	-	+	+
Dichloromethane fraction of <i>T. diversifolia</i> leaves.	+	-	+	+
Methanol-water fraction of <i>T. diversifolia</i> leaves.	-	-	+	+

The multiple regression analysis showed that total phenolic and flavonoid content contributed 92.1% and 90.6% of anti-inflammatory and antioxidant activity, respectively (Figure 2). The unstandardized predicted value was for anti-inflammatory (both from RBC membrane stabilization and protein denaturation inhibition method) and antioxidant activity. Total phenolic and flavonoid content significantly affected anti-inflammatory and antioxidant activity and were classified as highly correlated (Eryani et al., 2022).

Table 5. Total phenolic and flavonoid content of *T. diversifolia* leaves ethanolic extract and its fractions. *Description:* *Data was shown as mean \pm SD (n=3); different letter notations in the same parameter indicate significant differences (LSD, $p < 0.01$). GAE = gallic acid equivalent. QE = quercetin equivalent.

Sample	Total phenolic content* (mg GAE/g sample)	Total flavonoids content* (mg QE/g sample)
Ethanolic extract of <i>T. diversifolia</i> leaves	32.447 \pm 0.337 ^a	21.374 \pm 0.585 ^e
n-hexane fraction of <i>T. diversifolia</i> leaves	12.035 \pm 0.262 ^b	135.103 \pm 0.517 ^f
Dichloromethane fraction of <i>T. diversifolia</i> leaves	16.287 \pm 0.730 ^c	63.661 \pm 1.056 ^g
Methanol-water fraction of <i>T. diversifolia</i> leaves	26.349 \pm 0.330 ^d	11.901 \pm 0.119 ^h

Based on Pearson correlation analysis (Table 6), the correlation of total phenolic content to the anti-inflammatory activity (both from RBC membrane stabilization and protein denaturation inhibition method) was negative, meaning that the higher the phenolic content, the lower the IC₅₀ on RBC membrane stabilization and protein denaturation inhibition. The total phenolic content played a significant correlation to RBC membrane stabilization with the value of -0.768 (high negative correlation, $p < 0.01$), but no significant correlation to protein denaturation inhibition and antioxidant activity with a correlation coefficient of -0.626 (moderate negative correlation) and -0.403 (low negative correlation), respectively. The total flavonoid content negatively correlated to antioxidant activity, suggesting that the higher the total flavonoid content, the lower the IC₅₀ to antioxidant activity. On the other hand, the correlation between total flavonoid content and anti-inflammatory activity (RBC membrane stabilization and protein denaturation inhibition method) was positive. This anomaly is related to the anomaly in the total flavonoid content calculation due to the previously described challenges. There was no significant correlation between total flavonoid content and anti-inflammatory and antioxidant activity. The correlation coefficient on total flavonoid content to RBC membrane stabilization was 0.433 (low positive correlation), protein denaturation inhibition was 0.47 (low positive correlation), and antioxidant was -0.233 (negligible negative correlation) (Mukaka, 2012).





Figure 2. Multiple regression analysis showed that total phenolic (TPC) and flavonoid content (TFC) contributed significantly to the anti-inflammatory and antioxidant activity with high correlation.


These findings would complete the anti-inflammatory mechanism of action of ethanolic extract of *T. diversifolia* leaves and its fraction, following red blood cell membrane stabilization and protein denaturation inhibition. The antioxidant properties likely contributed more to RBC membrane stabilization than protein denaturation inhibition. The phenolic and flavonoid content contributed most to antioxidant and RBC membrane stabilization rather than protein denaturation inhibition (Figure 3).

Table 6. Pearson correlation analysis on total phenolic and flavonoid content to anti-inflammatory and antioxidant activity. *Description:* * correlation is significant at a 99% confidence level ($p < 0.01$).

Parameter	Pearson correlation value	
	Total phenolic content	Total flavonoid content
RBC stabilization membrane	-0.626	0.470
Protein denaturation inhibition	-0.768*	0.433
Antioxidant	-0.403	-0.801

The ethanolic extract of *T. diversifolia* leaves showed the highest anti-inflammatory and antioxidant activity. It contained terpenoids, alkaloids, polyphenols, and flavonoids. Terpenoids inhibit cyclooxygenase and suppress c-Jun N-terminal kinase, leading to the inhibition of pro-inflammatory mediators (Subedi & Yumnam, 2021).

Figure 3. A proposed mechanism on how the compounds contained in ethanolic extract of *T. diversifolia* leaves ethanolic extract and its fractions act as anti-inflammatory and antioxidant (Bouhlali et al., 2020; Katzung et al., 2018; Maleki et al., 2019; Souto et al., 2011; Subedi & Yumnam, 2021). *Description:* Reactive Oxygen Species (ROS),  (inhibit/block).



Alkaloids might inhibit either cyclooxygenase 1 or 2, as well as lipoxygenase, resulting in inhibition of pro-inflammatory mediators (Souto et al., 2011). Polyphenols stabilize membrane structure and function by forming stronger hydrogen bonds (Bouhlali et al., 2020). Mechanism of actions of flavonoids as anti-inflammatory and antioxidant agents are mediated by inhibition of regulatory proteins, e.g., protein kinases and phosphodiesterases, resulting in decreased signal transduction; scavenger activity on free radicals, resulting in decreased oxidative stress; inhibition of phospholipase A2, cyclooxygenase, and lipoxygenase, causing decrease release of pro-inflammatory mediators; modulation of transcription factors: NF- κ B, GATA-3, STAT-6, resulting in decreased transcription of pro-inflammatory genes; and also inhibition of cell activation, maturation, signaling transduction, secretory processes of immune cells causing immature immune cells, decreased cell proliferation, and decreased release of pro-inflammatory cytokines (Maleki et al., 2019). The possible mechanism of actions of compounds contained in

ethanolic extract of *T. diversifolia* leaves are shown in Figure 3. The fact that the ethanolic extract of *T. diversifolia* leaves had the highest anti-inflammatory and antioxidant activity compared to that of the fractions suggests that these compounds in the extract might exhibit synergistic additive effects, not antagonistic ones—the total phenolic and flavonoid concentrations in a plant correlate with the anti-inflammatory and antioxidant activity. The higher polyphenols lead to higher anti-inflammatory and antioxidant activity (Benjamin et al., 2024; Rini et al., 2023). Nevertheless, further research needs to be done to confirm these possibilities.

The ability to block or inhibit inflammation and the antioxidants featured by ethanolic extract of *T. diversifolia* leaves will act as a possible mechanism of the antiproliferative and anticancer activities of *T. diversifolia* leaves. Inflammation is one of the hallmarks of cancer (Hanahan, 2022). Thus, anti-inflammatory drugs have been repurposed for anticancer therapy (Ozleyen et al., 2023). On the other hand, the antioxidant properties of natural products will scavenge the free radicals in our body and control inflammatory responses, supporting the already considered anti-inflammatory activity. These two mechanisms of action, antioxidant and anti-inflammatory activities, may work synergistically in cancer chemoprevention (Jongrungrauchok et al., 2023).

4. CONCLUSION

Based on the results, we can conclude that the ethanolic extract of *T. diversifolia* leaves had the best anti-inflammatory properties compared to its fractions based on both methods and antioxidant activity. The antioxidant activity contributed more to RBC membrane stabilization compared to protein denaturation inhibition. The phenolic and flavonoid content contributed more to antioxidant and red blood cell membrane stabilization. The ethanolic extract of *T. diversifolia* leaves contained terpenoids, alkaloids, polyphenols, and flavonoids that might exert synergistic or additive effects on anti-inflammatory and antioxidant properties. Nevertheless, research is still needed to confirm these possibilities.

ACKNOWLEDGEMENT

The authors would like to thank the University of Jember for funding this research through Hibah KeRis 2022 and the students helping with data collection and analysis: Ahya Natasya, Anneke Putri Lestari Junaedi, Feny Saskia Safitri, Intan Nabila Sufi Zikrina, Shinta Dwi Kurniawati, and Ramadhan Bagus Cahyono.

CONFLICT OF INTEREST

All authors declared that there was no conflict of interest.

REFERENCES

- Benjamaa, R., Elbouny, H., Errati, H., Moujanni, A., Kaushik, N., Gupta, R., Ennibi, O. K., Nasser, B., Choi, E. H., Kaushik, N. K., & Essamadi, A. (2024). Comparative evaluation of antioxidant activity, total phenolic content, anti-inflammatory, and antibacterial

- potential of Euphorbia-derived functional products. *Frontiers in Pharmacology*, 15. <https://doi.org/10.3389/fphar.2024.1345340>
- Bindu, S., Mazumder, S., & Bandyopadhyay, U. (2020). Non-steroidal anti-inflammatory drugs (NSAIDs) and organ damage: A current perspective. In *Biochemical Pharmacology* (Vol. 180). Elsevier Inc. <https://doi.org/10.1016/j.bcp.2020.114147>
- Bouhlali, E. dine T., Hmidani, A., Bourkhis, B., Khouya, T., Ramchoun, M., Filali-Zegzouti, Y., & Alem, C. (2020). Phenolic profile and anti-inflammatory activity of four Moroccan date (*Phoenix dactylifera* L.) seed varieties. *Heliyon*, 6(2). <https://doi.org/10.1016/j.heliyon.2020.e03436>
- Bourais, I., Elmarrkechy, S., Taha, D., Mourabit, Y., Bouyahya, A., El Yadini, M., Machich, O., El Hajjaji, S., El Boury, H., Dakka, N., & Iba, N. (2023). A Review on Medicinal Uses, Nutritional Value, and Antimicrobial, Antioxidant, Anti-Inflammatory, Antidiabetic, and Anticancer Potential Related to Bioactive Compounds of *J. regia*. *Food Reviews International*. <https://doi.org/10.1080/87559129.2022.2094401>
- Christiana, R., Kristopo, H., & Limantara, L. (2008). Photodegradation and Antioxidant Activity of Chlorophyll a from *Spirulina* (*Spirulina* sp.) Powder. In *Indo. J. Chem* (Vol. 8, Issue 2).
- Dalimartha, S. (1999). *Atlas tumbuhan obat Indonesia*. Trubus Agriwidya.
- Eryani, I. G. A. P., Jayantari, M. W., & Wijaya, I. K. M. (2022). Sensitivity Analysis in Parameter Calibration of the WEAP Model for Integrated Water Resources Management in Unda Watershed. *Civil Engineering and Architecture*, 10(2). <https://doi.org/10.13189/cea.2022.100206>
- Greten, F. R., & Grivennikov, S. I. (2019). Inflammation and Cancer: Triggers, Mechanisms, and Consequences. In *Immunity* (Vol. 51, Issue 1, pp. 27–41). Cell Press. <https://doi.org/10.1016/j.immuni.2019.06.025>
- Hanahan, D. (2022). Hallmarks of Cancer: New Dimensions. In *Cancer Discovery* (Vol. 12, Issue 1, pp. 31–46). American Association for Cancer Research Inc. <https://doi.org/10.1158/2159-8290.CD-21-1059>
- Jongrungraungchok, S., Madaka, F., Wunnakup, T., Sudsai, T., Pongphaew, C., Songsak, T., & Pradubyat, N. (2023). In vitro antioxidant, anti-inflammatory, and anticancer activities of mixture Thai medicinal plants. *BMC Complementary Medicine and Therapies*, 23(1). <https://doi.org/10.1186/s12906-023-03862-8>
- Katzung, B. G., von Zastrow, M., Holford, N. H., Correia, M. A., Pappano, A. J., Biaggioni, I., Robertson, D., Benowitz, N. L., Harvey, R. D., Grant, A. O., Sam, R., Ives, H., Pearce, D., Reid, I. A., Hwa, J., & et al. (2018). *Basic & Clinical Pharmacology* (B. G. Katzung, Ed.; 14th Edition). McGraw-Hill Education.
- Lopes, D. C. D. X. P., De Oliveira, T. B., Viçosa, A. L., Valverde, S. S., & Ricci Júnior, E. (2021). Anti-Inflammatory Activity of the Compositae Family and Its Therapeutic Potential. In *Planta Medica* (Vol. 87, Issues 1–2, pp. 71–100). Georg Thieme Verlag. <https://doi.org/10.1055/a-1178-5158>
- López-Cabeza, R., Rodríguez-Sabina, S., Reyes, C. P., Expósito, D. G., Giménez, C., Jiménez, I. A., Cabrera, R., & Bazzocchi, I. L. (2024). Bio-guided isolation of aromatic abietane diterpenoids from *Salvia canariensis* as biopesticides in the control of phytopathogenic fungi. *Pest Management Science*, 80(4), 2199–2207. <https://doi.org/10.1002/ps.7958>
- Loucif, K., Benabdallah, H., Benchikh, F., Mehloos, S., Souici, C. Ben, & Amira, S. (2020). Total Phenolic Contents, DPPH Radical Scavenging and β -Carotene Bleaching Activities of Aqueous Extract from *Ammoides atlantica*. *Journal of Drug Delivery and Therapeutics*, 10(3-s), 196–198. <https://doi.org/10.22270/jddt.v10i3-s.4151>

- Lubis, B., Saputri, I. N., Ajartha, R., Bangun, S. M. B., Pranata, C., Purba, N., & Turnip, N. U. M. Br. (2020). Anti-inflammatory Activity Test for Ethanol Extract Moon Flower (*Tithonia diversifolia*) Leaves to Male White Mice. *Proceedings of the International Conference on Health Informatics and Medical Application Technology*, 551–557. <https://doi.org/10.5220/0009974705510557>
- Maleki, S. J., Crespo, J. F., & Cabanillas, B. (2019). Anti-inflammatory effects of flavonoids. In *Food Chemistry* (Vol. 299). Elsevier Ltd. <https://doi.org/10.1016/j.foodchem.2019.125124>
- Moulishankar, A. (2021). Significance of TLC and HPTLC in Phytochemical Screening of Herbal Drugs. In *Article in Journal of Global Pharma Technology*. www.jgpt.co.in
- Mukaka, M. M. (2012). Statistics Corner: A guide to appropriate use of Correlation coefficient in medical research. In *Malawi Medical Journal* (Vol. 24, Issue 3). www.mmj.medcol.mw
- Nasution, M.R., Ardhiyati, B., Tinggi, S., Riau, I. F., Kamboja, J., & Baru-Panam, S. (2019). *Total Fenolik dan Flavonoid serta Aktivitas Antioksidan Ekstrak Etanol Daun Tenggek Burung (Eudia redlevi)*.
- Nunes, C. dos R., Arantes, M. B., de Faria Pereira, S. M., da Cruz, L. L., de Souza Passos, M., de Moraes, L. P., Vieira, I. J. C., & de Oliveira, D. B. (2020). Plants as Sources of Anti-inflammatory Agents. In *Molecules* (Vol. 25, Issue 16). MDPI AG. <https://doi.org/10.3390/molecules25163726>
- Ozleyen, A., Yilmaz, Y. B., Donmez, S., Atalay, H. N., Antika, G., & Tumer, T. B. (2023). Looking at NSAIDs from a historical perspective and their current status in drug repurposing for cancer treatment and prevention. In *Journal of Cancer Research and Clinical Oncology* (Vol. 149, Issue 5, pp. 2095–2113). Springer Science and Business Media Deutschland GmbH. <https://doi.org/10.1007/s00432-022-04187-8>
- Phongpaichit, S., Nikom, J., Rungjindamai, N., Sakayaroj, J., Hutadilok-Towatana, N., Rukachaisirikul, V., & Kirtikara, K. (2007). Biological activities of extracts from endophytic fungi isolated from Garcinia plants. *FEMS Immunology and Medical Microbiology*, 51(3), 517–525. <https://doi.org/10.1111/j.1574-695X.2007.00331.x>
- Puspitasari, E., Ningsih, Y., Triatmoko, B., & Dianasari, D. (2022). Antiproliferative activity of ethanolic extract of kembang bulan (*Tithonia diversifolia*) leaf on HeLa cervical cancer cell line. *Indonesian Journal of Cancer Chemoprevention*, 13(1), 55–60.
- Rini, S., Shweta, V., Vijender, S., & Sakshi, B. (2023). Impact of geographical location on the polyphenolic content of *Nigella sativa* seed extract and the relative anti-inflammatory and antioxidant potential thereof: A comparative study. *Indian Journal of Natural Products and Resources*, 14(1), 89–96. <https://doi.org/10.56042/ijnpr.v14i1.1132>
- Saleem, A., Saleem, M., & Akhtar, M. F. (2020). Antioxidant, anti-inflammatory and antiarthritic potential of *Moringa oleifera* Lam: An ethnomedicinal plant of Moringaceae family. *South African Journal of Botany*, 128, 246–256. <https://doi.org/10.1016/j.sajb.2019.11.023>
- Satria, R., Hakim, A. R., & Darsono, P. V. (2022). Penetapan Kadar Flavonoid Total Dari Fraksi n-Heksana Ekstrak Daun Gelinggang dengan Metode Spektrofotometri UV-Vis. *Journal of Engineering, Technology, and Applied Science*, 4(1), 33–46. <https://doi.org/10.36079/lamintang.jetas-0401.353>
- Siegel, R. L., Miller, K. D., Wagle, N. S., & Jemal, A. (2023). Cancer statistics, 2023. *CA: A Cancer Journal for Clinicians*, 73(1), 17–48. <https://doi.org/10.3322/caac.21763>
- Souto, A. L., Tavares, J. F., Da Silva, M. S., De Diniz, M. F. F. M., De Athayde-Filho, P. F., & Barbosa Filho, J. M. (2011). Anti-inflammatory activity of alkaloids: An update from 2000 to 2010. In *Molecules* (Vol. 16, Issue 10, pp. 8515–8534). <https://doi.org/10.3390/molecules16108515>

- Subedi, L., & Yumnam, S. (2021). Terpenoids from abies holophylla attenuate lps-induced neuroinflammation in microglial cells by suppressing the jnk-related signaling pathway. *International Journal of Molecular Sciences*, 22(2), 1–12. <https://doi.org/10.3390/ijms22020965>
- Tamfu, A. N., Roland, N., Mfifen, A. M., Kucukaydin, S., Gaye, M., Botezatu, A. V., Emin Duru, M., & Dinica, R. M. (2022). Phenolic composition, antioxidant and enzyme inhibitory activities of Parkia biglobosa (Jacq.) Benth., Tithonia diversifolia (Hemsl) A. Gray, and Crossopteryx febrifuga (Afzel.) Benth. *Arabian Journal of Chemistry*, 15(4). <https://doi.org/10.1016/j.arabjc.2021.103675>
- Tu, Z., Zhong, Y., Hu, H., Shao, D., Haag, R., Schirner, M., Lee, J., Sullenger, B., & Leong, K. W. (2022). Design of therapeutic biomaterials to control inflammation. In *Nature Reviews Materials* (Vol. 7, Issue 7, pp. 557–574). Nature Research. <https://doi.org/10.1038/s41578-022-00426-z>
- World Health Organization. (2022). *WHO Statistics 2022: Monitoring Health for the SDGs*. <http://apps.who.int/bookorders>.
- Xin, Y. J., Choi, S., Roh, K. B., Cho, E., Ji, H., Weon, J. B., Park, D., Whang, W. K., & Jung, E. (2021). Anti-inflammatory activity and mechanism of isookanin, isolated by bioassay-guided fractionation from bidens pilosa l. *Molecules*, 26(2). <https://doi.org/10.3390/molecules26020255>
- Yesmin, S., Paul, A., Naz, T., Rahman, A. B. M. A., Akhter, S. F., Wahed, M. I. I., Emran, T. Bin, & Siddiqui, S. A. (2020). Membrane stabilization as a mechanism of the anti-inflammatory activity of ethanolic root extract of Choi (Piper chaba). *Clinical Phytoscience*, 6(1). <https://doi.org/10.1186/s40816-020-00207-7>
- Yu, X., Yang, T., Qi, Q., Du, Y., Shi, J., Liu, X., Liu, Y., Zhang, H., Zhang, Z., & Yan, N. (2021). Comparison of the contents of phenolic compounds including flavonoids and antioxidant activity of rice (Oryza sativa) and Chinese wild rice (Zizania latifolia). *Food Chemistry*, 344. <https://doi.org/10.1016/j.foodchem.2020.128600>
- Zhang, N., Wang, M., Li, Y., Zhou, M., Wu, T., & Cheng, Z. (2021). TLC–MS identification of alkaloids in Leonuri Herba and Leonuri Fructus aided by a newly developed universal derivatisation reagent optimised by the response surface method. *Phytochemical Analysis*, 32(3), 242–251. <https://doi.org/10.1002/pca.2970>
- Zhao, G. J., Xi, Z. X., Chen, W. S., Li, X., Sun, L., & Sun, L. N. (2012). Chemical constituents from Tithonia diversifolia and their chemotaxonomic significance. *Biochemical Systematics and Ecology*, 44, 250–254. <https://doi.org/10.1016/j.bse.2012.06.019>
- Zulkifli, S. A., Gani, S. S. A., Zaidan, U. H., & Halmi, M. I. E. (2020). Optimization of Total Phenolic and Flavonoid Contents of Defatted Pitaya (Hylocereus polyrhizus) Seed Extract and Its Antioxidant Properties. *Molecules*, 25(4). <https://doi.org/10.3390/molecules25040787>